

illness newly admitted to nursing homes. *Psychiatric Services* (Washington, D.C.) 62:1098–1100, 2011

54. Lan CC, Su TP, Chen YS, et al: Treatment dilemma in comorbidity of schizophrenia and idiopathic Parkinson's disease. *General Hospital Psychiatry* 33:411e3–411e5, 2011
55. Hus JH, Chien IC, Lin CH, et al: Incidence of diabetes in patients with schizophrenia: a population-based study. *Canadian Journal of Psychiatry* 56:19–26, 2011
56. Felmet K, Zisook S, Kasckow JW: Elderly patients with schizophrenia and depression:

diagnosis and treatment. *Clinical Schizophrenia and Related Psychoses* 4:239–250, 2011

57. Guo JJ, Keck PE, Li H, et al: Diabetes associated with antipsychotic use among Medicaid patients with bipolar disorders: a nested case-control study. *Pharmacotherapy* 27:27–35, 2007
58. Howe AM, Kozma C, Russo P, et al: The prevalence of potential antipsychotic drug-drug interactions in a large US national managed care database. Presented at the Annual US Psychiatric and Mental Health Congress, New Orleans, La, Nov 15–19, 2006


59. Frois C, Guerin A, Saraogi A, et al: Perceptions and prescribing considerations among US psychiatrists regarding drug-drug interactions associated with oral atypical antipsychotics. *Current Medical Research and Opinion* 26:2735–2744, 2010
60. Cavuto NJ, Woosley RL, Sale M: Pharmacies and prevention of potentially fatal drug interactions. *JAMA* 275:1086–1087, 1996
61. Dorsey ER, Rabbani A, Gallagher SA, et al: Impact of FDA black box advisory on antipsychotic medication use. *Archives of Internal Medicine* 170:96–103, 2010

Correction to Bassilios et al., 2012

In the article “Enhanced Primary Mental Health Services in Response to Disaster,” by Bridget Bassilios, D.Psych., Lennart Reifels, Dipl.-Psych., and Jane Pirkis, Ph.D. (September 2012 issue, pp. 868–874), the legend of Figure 1, on page 870, is incorrect. The key should indicate that the curve represents sessions and the bars represent referrals. A corrected figure appears below.

Figure 1

Monthly referrals for mental health care through Australia's Access to Allied Psychological Services program after the 2009 bushfires, January 2009 to June 2011^a



^a An additional five referrals and ten sessions were recorded as having taken place pre-January 2009, but the dates are likely to be data entry errors.