- 6. Institute of Medicine: Improving the Quality of Mental Health Care for Mental And Substance-Use Conditions. Washington, DC, National Academies Press, 2006
- McLellan AT, Starrels JL, Tai B, et al: Can substance use disorders be managed using the chronic care model? Review and recommendations from a NIDA consensus group. Public Health Reviews 35:1–14, 2014
- Moyer VA: Screening and behavioral counseling interventions in primary care to reduce alcohol misuse: US Preventive Services Task Force recommendation statement. Annals of Internal Medicine 159:210–218, 2013
- Gardner EM, McLees MP, Steiner JF, et al: The spectrum of engagement in HIV care and its relevance to test-and-treat strategies for prevention of HIV infection. Clinical Infectious Diseases 52: 793–800, 2011
- Saitz R: Lost in translation: the perils of implementing alcohol brief intervention when there are gaps in evidence and its interpretation. Addiction 109:1060–1062, 2014
- Saitz R, Cheng DM, Winter M, et al: Chronic care management for dependence on alcohol and other drugs: the AHEAD randomized trial. JAMA 310:1156–1167, 2013
- 12. Oslin DW, Lynch KG, Maisto SA, et al: A randomized clinical trial of alcohol care management delivered in Department of Veterans

Affairs primary care clinics versus specialty addiction treatment. Journal of General Internal Medicine 29:162–168, 2014

- Weisner C, Mertens J, Parthasarathy S, et al: Integrating primary medical care with addiction treatment: a randomized controlled trial. JAMA 286:1715–1723, 2001
- 14. Tai B, McLellan AT: Integrating information on substance use disorders into electronic health record systems. Journal of Substance Abuse Treatment 43:12–19, 2012
- Smith SA, Shah ND, Bryant SC, et al: Chronic care model and shared care in diabetes: randomized trial of an electronic decision support system. Mayo Clinic Proceedings 83:747–757, 2008
- 16. Otte-Trojel T, de Bont A, van de Klundert J, et al: Characteristics of patient portals developed in the context of health information exchanges: early policy effects of incentives in the meaningful use program in the United States. Journal of Medical Internet Research 16:e258, 2014
- Mauer BJ: Behavioral Health/Primary Care Integration: Finance, Policy and Integration of Services. Washington, DC, National Council for Community Behavioral Healthcare, 2006
- Langley G, Moen R, Nolan K, et al: The Improvement Guide: A Practical Approach to Enhancing Organizational Performance. San Francisco, Jossey-Bass, 2009

Correction to Wiegand and Godemann (2017)

The Brief Report "Increased Treatment Complexity for Major Depressive Disorder for Inpatients With Comorbid Personality Disorder," by Hauke F. Wiegand, M.D., Ph.D., and Frank Godemann, M.D., published online in *Psychiatric Services* in Advance on February 1, 2017, contained an error. In Table 1, the values for males and females are reversed. For major depression, the correct values are 24,311 females and 17,245 males. For major depression + personality disorder, the correct values are 5,704 females and 3,026 males. A corrected table will appear in the May issue (*Psychiatric Services* vol. 68, no. 5), and a corrected version was uploaded to *Psychiatric Services* in Advance on March 8, 2017.