

# A Randomized Controlled Trial of a Money Management–Based Substance Use Intervention

Marc I. Rosen, M.D.

Kathleen M. Carroll, Ph.D.

Elina Stefanovics, Ph.D.

Robert A. Rosenheck, M.D.

**Objective:** Money management has been implemented, often in bundled interventions, as a strategy to counteract spending of public support checks and other funds on drugs and alcohol. The authors conducted a randomized controlled trial of a voluntary money management program as an adjunctive treatment for patients in treatment for mental illness, substance use disorders, or both. In the advisor-teller money manager (ATM) intervention, a money manager stores patients' checkbooks and automated bank cards, trains patients to manage their own funds, and links spending to activities related to their treatment goals. **Methods:** Eighty-five veterans with recent use of alcohol or cocaine were randomly assigned to 36 weeks of the ATM intervention or a control intervention (completion of a simple financial workbook). **Results:** With ATM, 75% of veterans gave their checkbook to their money manager to hold, and participants attended significantly more therapy sessions than those assigned to the control therapy (mean of 20.6 versus 8.1 sessions). Although participants assigned to ATM did not show significantly greater improvement over time on the primary outcomes (self-reported abstinence from alcohol and cocaine and negative urine tests for cocaine metabolite), they reduced their Addiction Severity Index drug and alcohol use composite scale scores more rapidly than the control group. High rates of abstinence in both groups created a ceiling effect, limiting the power to detect improved abstinence rates. **Conclusions:** In this relatively small trial, ATM, a money management intervention, showed promise in engaging patients, improving their money management, and improving some substance abuse outcomes. (*Psychiatric Services* 60:498–504, 2009)

Substance abuse is common among people with serious mental illness (1), many of whom receive disability payments. People prone to addiction show higher use of drugs and alcohol when checks are received at the be-

ginning of the month (2,3). Harmful consequences of drug use that also occur disproportionately when checks are received include discharge from residential substance abuse treatment (4), emergency services visits (5), and deaths related to drug and

alcohol use (6). Money itself can be a relapse trigger (7), and benefit checks can facilitate the purchase of drugs and alcohol.

Based on the connection between substance use and mismanagement of personal funds, many treatment programs have instituted efforts to help psychiatric patients with co-occurring psychiatric and substance abuse problems to avoid diverting funds toward substance use (8,9). Money management services range from voluntary programs designed to provide information about budgeting (10) to involuntary, mandated interventions, such as the assignment of a representative payee to receive and manage a patient's check. Assertive community treatment programs that treat patients with co-occurring substance use and psychiatric disorders often help with budgeting, storing funds, paying bills, and managing entitlements, but money management is only one of several interventions in assertive community treatment (11–13), making it difficult to pinpoint the effects of money management among the other services.

In the sole randomized controlled trial involving involuntary money management, 184 veterans were randomly assigned to referral to a community-based payee program or to treatment as usual and followed for 12 months (14). Compared with beneficiaries in the control group, beneficiaries in the payee assignment group used substances for fewer months and rated their money management

---

*The authors are affiliated with the Department of Veterans Affairs (VA) Connecticut Healthcare System, West Haven. Dr. Rosen, Dr. Carroll, and Dr. Rosenheck are also with the Department of Psychiatry, 116A, Yale University, West Haven, CT 06525 (e-mail: marc.rosen@yale.edu).*

as significantly better, but this pioneering trial's generalizability was limited by the fact that the active intervention was bundled with and included referral to extensive case management services in addition to the payee assignment. The benefits of payee assignment have been equivocal in other studies (3,15), although specialized payee services that are based at community mental health centers have shown promise (16).

There have not been any controlled trials of an unbundled money management-based intervention. In this article we report on a 36-week single-blind, randomized controlled clinical trial designed to test the efficacy of a voluntary money management program, which we compared with a control condition, in fostering abstinence among veterans in treatment for mental illness, a substance use disorder, or co-occurring disorders. Secondary aims of the study were to determine to what extent participants engaged in money management—that is, attending sessions and allowing their funds to be managed—and how participants rated the experience.

## Methods

### *Recruitment, enrollment, and randomization*

Patients who were being treated for a psychiatric disorder, a substance use disorder, or co-occurring psychiatric and substance use disorders were recruited by advertisement and clinician referral at the main campuses of the Veterans Affairs (VA) Connecticut Healthcare System and the Northampton VA Medical Center. Patients were enrolled who had spent at least \$100 on cocaine, alcohol, cannabis, or a combination of these substances during the preceding 90 days and received at least \$300 income per month. Patients were excluded if they already had a payee or other money manager or if they were physiologically dependent on opioids or alcohol. All study procedures were approved by the institutional review boards of the VA Connecticut Healthcare System and Yale University. All patients provided written informed consent, and enrollment was conducted from August 2000 until February 2004.

Patients were screened to deter-

mine eligibility, and eligible patients completed an extensive battery of assessments. Before we randomly assigned participants to an intervention condition, all patients were scheduled for an individual visit with the money manager to assess their finances and to explain the interventions in more detail. To avoid assigning patients who had no interest in money management to an intervention, random assignment occurred only for patients who came to the visit with funds to open a checking account or with their checkbook so that they could begin receiving advisor-teller money management (ATM), should they be assigned to that intervention. The substance for which each patient had spent the most in the previous 90 days was balanced between the ATM and control groups by urn randomization (17).

### *Interventions*

**ATM.** The ATM intervention is a therapy that uses voluntary money management to reduce substance use and incorporates elements of optimal payee practice (18). Staff trained as money managers in this program performed three functions—serving as a teller to store patients' funds, training patients to manage their own funds, and linking spending to treatment goals. In the teller function, staff encouraged patients, but did not require them, to establish a checking account and to allow the money manager to store their checkbook, their automated bank card, or both. In the training function, patients and counselors planned a budget each month, and at the end of the month, they reviewed whether the budget had been followed.

To encourage treatment-linked spending, breath-analysis tests and on-site urine tests were conducted at weekly ATM counseling visits. Patients were encouraged not to spend money if the tests were positive. Patients were encouraged to identify additional activities compatible with abstinence (19) and make contracts to reward themselves the next week with their own discretionary funds if the activities were successfully carried out or if they had been abstinent. Patients who requested funds that had not been budgeted were encouraged to fill out a

written request and wait two working days to be certain that the unplanned expenditure was prudent. An updated manual describing ATM is available for download (20).

**Control.** For the control condition, patients were given a workbook in which they were asked to construct a budget and track their expenses and income. Counseling sessions involved supporting patients' efforts to use the workbook. The same bachelor's-level staff who served as ATM managers also guided the control intervention.

### *Data collection and measurement*

Participants were paid for each of the 19 assessments and could earn a total of \$325 if they completed all of the study assessments. Assessments were conducted by bachelor's-level staff. Baseline assessments characterized patients on a broad array of measures. Interviews using the Structured Clinical Interview for DSM-IV (21) were conducted, and the Addiction Severity Index (ASI) (22) was administered.

Breath-analysis tests, urine toxicology tests, and substance abuse calendars, completed with the timeline follow-back method (23), were collected every two weeks. On-site toxicology tests (Dade Behring, San Jose) were administered and processed for cocaine, opioids, and cannabis. The follow-up version of the ASI was administered every four weeks. Observer-rated psychosocial function was measured every four weeks with the Global Assessment of Functioning (GAF) (24).

Domains of money management were assessed every four weeks by having patients rate their agreement with statements rated on a Likert scale anchored by 1, strongly disagree, and 4, strongly agree (items available on request). The domains assessed, drawn from prior research (25), were budgeting, perceived benefit from the intervention, degree of misspending, satisfaction with the intervention, and coercion by the money manager.

### *Data analysis*

Data analysis proceeded in several steps. First, baseline characteristics of participants assigned to ATM and those assigned to the control group

were compared by *t* tests and chi square tests to evaluate the success of the randomization and to identify unequally distributed characteristics to use later as covariates when comparing outcomes between the groups. To assess differences in the completeness of outcome data collected in the ATM and control groups, we compared time to last data collection by proportional hazards regression.

The primary outcome measures were weeks of self-reported abstinence from both alcohol and cocaine and proportion of urine toxicology tests free of cocaine metabolite. Secondary outcome measures considered included weeks of self-reported abstinence from cocaine, self-reported abstinence from alcohol, composite follow-up scores on the ASI alcohol and drug subscales (26), the amount of money spent on drugs and alcohol, and GAF scores.

Generalized estimation equations with SAS PROC GENMOD were used for the dichotomous variables representing self-reported abstinence and toxicology-verified abstinence. Hierarchical linear models using PROC MIXED were used to compare the effects of group assignment on continuous outcomes, such as the follow-up scores on the ASI composite indices.

## Results

### *Participants and sampling*

Altogether, 167 of the 391 individuals screened (43%) did not meet eligibility criteria. Fifty-nine percent (133 veterans) of the 224 eligible participants were not assigned to an intervention; 36% (81 veterans) refused to participate, and an additional 23% (52 veterans) were excluded for other reasons (for example, they did not keep appointments or moved out of the geographic area). Eighty-five patients were randomly assigned to an intervention (44 to ATM and 41 to the control condition) and completed at least one follow-up assessment.

Forty of the 85 patients who were followed completed week 36 assessments, 20 of whom were assigned to ATM and 20 to the control condition. There was no significant difference between groups in length of time in which data were collected,

and there was no difference in the mean $\pm$ SD percentage of scheduled monthly follow-up assessments completed (control, 65% $\pm$ 34%; ATM, 58% $\pm$ 33%). Three study participants died from medical illnesses, two in ATM and one assigned to the control group.

### *Participant characteristics*

Participants were predominantly middle-aged, male veterans (Table 1). Median monthly income was \$655, and median indebtedness was remarkably high (\$3,997). Seventy (82%) of the veterans owed more than \$100.

A substantial proportion of the participants had concomitant psychiatric illnesses. Altogether, 28 (33%) met *DSM-IV* criteria for major depressive disorder, four (5%) for schizophrenia or an unspecified psychotic disorder, ten (12%) for bipolar disorder, and five (6%) for generalized anxiety disorder or dysthymic disorder. Altogether, 52 (61%) of the participants had one of these psychiatric disorders.

Two of 27 baseline measures suggested more severe substance use in the ATM group than in the control group. Participants assigned to ATM had significantly higher baseline ASI alcohol composite scores ( $p=.01$ ) and had been hospitalized more times for substance abuse treatment ( $p=.03$ ) than participants in the control group.

### *Extent of participation in ATM*

As expected, compared with participants assigned to the control intervention, participants assigned to ATM attended many more money management sessions (20.6 $\pm$ 14.9 versus 8.1 $\pm$ 8.9 sessions;  $t=4.7$ ,  $df=71$ ,  $p<.001$ ). Altogether, 22 of the 44 participants assigned to ATM attended more than 18 sessions. Twenty-seven (61%) of those assigned to ATM gave checkbooks to their money managers to hold, and an additional six (14%) gave the money manager a checkbook to hold but took it back later.

Weekly treatment contracting was encouraged but was not used extensively. Twenty of 44 participants (46%) assigned to ATM rewarded themselves with their own funds (via check or automated bank card) for

completing activities they had contracted, but these participants reinforced themselves only an average of 2.6 times each. Reinforcing involved the veteran's writing a check to pay an expense or borrowing the stored bank card to make a withdrawal. The agreement to delay requests of unbudgeted funds to allow reconsideration of impulsive spending was rarely implemented; only five participants (11%) agreed to a delay of a day or longer before receiving requested funds.

### *Substance abuse outcomes*

The proportion of outcome observations with self-reported abstinence from both cocaine and alcohol was 86% in the control group and 88% in the ATM group (Table 2). Analysis yielded no significant differences associated with group assignment, time, or the group-by-time interaction (Table 2). The other primary outcome, proportion of toxicology tests negative for cocaine, was high (control, 88%; ATM, 92%) and did not differ significantly between groups overall or over time. However, a post hoc analysis restricted to 37 patients who had primarily abused cocaine supported the efficacy of ATM; after covarying for baseline measurements, there were significantly more negative toxicology tests in the ATM group ( $p=.008$ ). Results of analyses that separately considered self-reported abstinence from alcohol and abstinence from cocaine did not differ by group assignment (Table 2).

Table 3 shows the continuous outcome measures that were analyzed with mixed models. Study participants had less severe ASI drug composite scores over time, and this reduction in the drug composite scores was greater over time among ATM participants, as indicated by a significantly positive group-by-time effect. A similar pattern of results was apparent for the ASI alcohol composite scores. There was no difference between the groups by time in the amount of money spent on drugs and alcohol.

### *Money management processes*

Compared with participants in the control group, participants in the ATM intervention rated higher agree-

**Table 1**

Baseline characteristics of 85 veterans randomly assigned to the advisor-teller money management intervention or to a control intervention

Characteristic	Control (N=41)		Advisor-teller money manager (N=44)		Test statistic	df
	N	% <sup>a</sup>	N	% <sup>a</sup>		
Demographic						
Age (M±SD)	45.7±7.7		46.6±6.9		t=−.56	83
Gender (male)	40	98	44	100	χ <sup>2</sup> =1.09	1
Caucasian	16	40	26	61	χ <sup>2</sup> =3.47	1
Never married	19	46	17	39	χ <sup>2</sup> =.52	1
Living with significant other	15	38	15	34	χ <sup>2</sup> =.11	1
High school graduate	37	90	42	96	χ <sup>2</sup> =.88	1
Worked for pay in the past 30 days	18	45	17	39	χ <sup>2</sup> =.35	1
In a controlled environment in the past 30 days	19	48	19	48	χ <sup>2</sup> =.00	1
Days out of 30 in a controlled environment (M±SD)	13.7±19		7.4±8.4		t=1.53	52
Monthly income (M±SD \$)	631±565		711±624		t=−.60	78
Receiving SSI or SSDI <sup>b</sup>	9	22	9	21	χ <sup>2</sup> =.03	1
Receiving veterans benefits	11	27	10	23	χ <sup>2</sup> =.19	1
Receiving Social Security or veterans benefits	19	46	16	36	χ <sup>2</sup> =.87	1
Amount owed (M±SD \$)	8,881±16,570		8,763±11,950		t=.04	83
Substance abuse						
Primary substance was cocaine	21	51	16	36	χ <sup>2</sup> =1.90	1
Used alcohol in prior 30 days	17	42	17	39	χ <sup>2</sup> =.07	1
Used cocaine in prior 30 days	11	27	14	32	χ <sup>2</sup> =.25	1
Used cannabis in prior 30 days	4	10	3	7	χ <sup>2</sup> =.24	1
Days used primary drug in the past 30	2.6±4.3		3.8±5.8		t=−1.09	83
Positive toxicology for any substance	10	25	9	21	χ <sup>2</sup> =.19	1
Alcohol composite ASI score <sup>c</sup>	.24±.20		.36±.24		t=−2.50**	83
Drug composite ASI score <sup>c</sup>	.10±.07		.11±.11		t=−.32	83
Number of prior substance abuse treatments (M±SD)	9.5±9.6		18.1±23.6		t=−2.23*	58
Psychiatric comorbidity						
Ever hospitalized for a psychiatric problem	18	45	25	57	χ <sup>2</sup> =1.17	1
Number of psychiatric admissions	3.1±6.5		1.4±1.9		t=1.68	82
Received disability pay for psychiatric illness	9	26	7	16	χ <sup>2</sup> =1.16	1
Global Assessment of Functioning score (M±SD) <sup>d</sup>	55.4±12.6		53.4±9.6		t=1.55	77

<sup>a</sup> The baseline measures were collected at two timepoints and slightly varied at the two times.

<sup>b</sup> SSI, Supplemental Security Income; SSDI, Social Security Disability Insurance

<sup>c</sup> ASI, Addiction Severity Index. Possible scores range from 0 to 1, with higher scores indicating more severe problems.

<sup>d</sup> Possible scores range from 0 to 100, with higher scores indicating better functioning.

\*p=.03

\*\*p=.01

ment with statements about having benefited from treatment and agreed more with statements indicating that they planned and adhered to a budget (Table 4). These rated advantages of ATM did not appear to be related to nonspecific appreciation of ATM because overall satisfaction ratings did not differ between the two groups. Ratings of coercion were low and did not differ between groups, and the trend toward a group-by-time effect on coercion was not in the direction of more coercion over time in the ATM group.

### *Adverse effects of ATM*

One participant in the ATM intervention suspected that his money manager was responsible for his being billed by the VA for his appointments. A participant with schizophrenia temporarily believed that the VA had taken his money but changed his mind with treatment. Another participant with a history of irritability became upset during counseling and discontinued study participation. There were no other adverse events related to the money management intervention.

### *GAF*

There were no significant changes in GAF scores over time and no significant differences between groups (Table 3).

### **Discussion and conclusions**

ATM was well received, in that a substantial proportion of people assigned to this voluntary intervention attended money management sessions, stored their checkbooks or bank cards with the money manager, and contracted to receive more of their funds when they achieved individually cho-



**Table 2**

Generalized estimating equation analyses of self-reported abstinence and toxicology-verified abstinence for 85 veterans assigned to a money manager or to a control intervention<sup>a</sup>

Outcome	Abstinence (%)													
	Control (N=41)		Money manager (N=44)		Group			Week			Group × week			N <sup>b</sup>
	M	SD	M	SD	Estimate	z	p	Estimate	z	p	Estimate	z	p	
Primary														
Self-reported abstinence from alcohol and cocaine	86	35	88	32	−.34	−.63	ns	−.03	−1.56	ns	.01	.38	ns	2,032
Negative toxicology for cocaine	88	32	92	28	.20	.36	ns	−.03	−3.69	<.001	−.02	−1.00	ns	871
Secondary														
Self-reported abstinence from alcohol	88	32	88	32	−.06	−.12	ns	−.03	−1.61	ns	.02	.79	ns	2,034
Self-reported abstinence from cocaine	95	22	98	15	−.15	−.17	ns	−.002	−.05	ns	−.05	−1.05	ns	2,032

<sup>a</sup> Higher (more positive) group coefficients of abstinence measures indicate higher proportions abstinent in the control group.

<sup>b</sup> Number of observations used in the analysis

sen goals. Despite a ceiling effect caused by the overall high levels of abstinence among veterans who were followed, ATM showed promise as a treatment.

ATM was not associated with significant benefit, as shown by a group-by-time interaction, on self-reported abstinence or on cocaine-negative urine toxicology tests, but the intervention was associated with significant reductions in the ASI alcohol and drug abuse composite scores. Unlike dichotomous meas-

ures of abstinence or use, ASI composite scores are continuous measures reflecting other factors, such as the extent of drug-related problems and how much treatment is needed. One possible explanation for the effect on the drug and alcohol use composite scores and lack of significant effects on self-report and toxicology is that ATM may mitigate the severity of substance-related problems without reducing the proportion of weeks of substance use. It is noteworthy that the patterns of

group-by-time effects concerning cocaine use all were in the direction of efficacy of the ATM intervention. In addition to the significant effect on drug use composite scores among participants assigned to ATM, there were statistically nonsignificant trends toward less self-reported cocaine use, fewer positive urine toxicology tests, and less money spent on substances of abuse.

A money management intervention was unacceptable to many potential participants, as suggested by

**Table 3**

Mixed-models analyses of continuous outcome measures for 85 veterans assigned to a money manager or to a control intervention

Outcome	Control <sup>a</sup> (N=41)		Money manager (N=44)		Group			Week			Group × week			N <sup>b</sup>
	M	SD	M	SD	Estimate	z	p	Estimate	z	p	Estimate	z	p	
Addiction Severity Index follow-up score <sup>c</sup>														
Drug composite	.07	.07	.06	.07	-.004	-.46	ns	-.002	-4.86	<.001	.001	2.86	.004	489
Alcohol composite	.18	.17	.24	.18	-.05	-2.15	.03	-.003	-3.60	<.001	.003	2.44	.015	489
Spent in past 30 days on drugs and alcohol (\$)	46.80	162.41	60.15	212.29	-64.4	-1.69	.09	-5.02	-3.39	<.001	3.30	1.58	ns	489
Global Assessment of Functioning score <sup>d</sup>	58.4	12.4	57.4	12.4	-.40	-.26	ns	.10	1.59	ns	.04	.44	ns	457

<sup>a</sup> Percentage of ratings during follow-up period by control group participants without use of substance indicated

<sup>b</sup> Number of observations used in the analysis

<sup>c</sup> Higher (more positive) group coefficients of composite measures indicate more severe problems in the control group. Possible composite scores range from 0 to 1, with higher values indicating more severe problems.

<sup>d</sup> Possible scores range from 0 to 100, with higher scores indicating better functioning.

**Table 4**

Mixed-models analyses of money management domains for 85 veterans assigned to a money manager or to a control intervention<sup>a</sup>

Domain	Likert score <sup>b</sup>													N <sup>c</sup>
	Control (N=41)		Money manager (N=44)		Group			Week			Group × week			
	M	SD	M	SD	Estimate	z	p	Estimate	z	p	Estimate	z	p	
Budgeting	3.2	.8	3.6	.6	−.44	−2.6	.01	.001	.1	ns	.002	.2	ns	177
Perceived benefit	3.3	.8	3.7	.4	−.39	−2.6	.01	.006	1.1	ns	.000	.03	ns	175
Misspending	2.2	1.0	2.1	1.0	.24	1.0	ns	.01	1.3	ns	−.01	−.74	ns	176
Satisfaction	3.7	.7	3.8	.5	−.26	1.9	.06	.008	1.5	ns	.007	.9	ns	174
Coercion	1.4	.9	1.4	.8	−.28	−1.3	ns	−.02	2.0	.052	.02	2.0	.052	164

<sup>a</sup> Positive group coefficients indicate higher values in the control group.

<sup>b</sup> Possible scores on items for patients' assessment of the domains range from 1, strongly disagree, to 4, strongly agree.

<sup>c</sup> Number of observations used in the analysis

the fact that only 41% of eligible participants proceeded to the random assignment portion of the trial, but it was impressive that participation in ATM was extensive after randomization and that subjective ratings of ATM were quite favorable. These results are consistent with those of other studies in which participants in a community-based payee program indicated that although enrollment was involuntary, patients found money management helpful over time (27). Participation in ATM also reflected the fact that the participants wanted help with their finances. The desire for help with finances was also reflected in the surprisingly high participation in a control condition that offered little counselor input beyond support for completing a budgeting workbook. People with little income and people who have serious mental illness indicate a desire for help with concrete financial concerns (28), and the satisfaction of material needs typically takes precedence over other needs (29).

The tested version of advisor-teller money management involved storing checkbooks and bank cards instead of a version of ATM more akin to representative payee practice, in which the money manager receives the patient's funds and stores them in an account that can be accessed only by the payee. Although most veterans assigned to ATM agreed to have their savings materials stored and to meet a money

manager to budget, few participants (five out of 44 ATM participants) allowed the money manager to delay unplanned spending to reconsider the unplanned expenditure. It is possible that the intervention's effects would be greater if the budgeting and training components of ATM were used in conjunction with the restrictions on beneficiary access to funds implemented by assigned representative payees.

Limitations of this study include the restriction to veterans, almost all of whom were male, and high abstinence rates in both groups. Several factors mitigated against finding a robust effect of ATM in this population. The high rates of abstinence among study participants created a control group ceiling effect that was hard to improve on. This ceiling effect did not appear to be caused by selective dropout of participants with more severe baseline addiction, because post hoc analyses showed no association between baseline addiction severity measures and percentage of follow-up interviews that were completed. Further evidence that selective attrition did not account for the high abstinence rates comes from another study of patients with psychiatric disorders and concomitant substance abuse, conducted at the same sites and during the same period as our study, which also reported extremely high abstinence rates (30). Our randomly assigned sample of 85 had suf-

ficient power to detect an effect size of .4 with power of .8 and significance of .05, two-tailed (31), but enrollment of people who used alcohol alone also may have diluted the study's power to detect an effect of ATM on cocaine use. Although the "check effect" of more substance use when checks are received at the beginning of the month has been demonstrated for both alcohol and other drug use (6), approaches based on money management may be more effective for cocaine because cocaine binges are more expensive.

In summary, this study found that a money management-based treatment targeting cocaine and alcohol use is feasible, acceptable to participants, and potentially efficacious.

### Acknowledgments and disclosures

This research was supported in part by VA Research Development; by grants R01-DA012952, K02-DA017277, and P50-DA09241 from the National Institute on Drug Abuse; and by Veterans Integrated Service Network 1 of the Mental Illness Research, Education and Clinical Care Center.

The authors report no competing interests.

### References

1. Regier DA, Farmer ME, Rae DS, et al: Comorbidity of mental disorders with alcohol and other drug abuse: results from the Epidemiologic Catchment Area (ECA) Study. *JAMA* 264:2511-2518, 1990
2. Shaner A, Eckman TA, Roberts LJ, et al: Disability income, cocaine use, and repeated hospitalization among schizophrenic cocaine abusers: a government-sponsored re-

- volving door. *New England Journal of Medicine* 333:777–783, 1995
3. Swartz JA, Hsieh CM, Baumohl J: Disability payments, drug use and representative payees: an analysis of the relationships. *Addiction* 98:965–975, 2003
4. Satel S, Reuter P, Hartley D, et al: Influence of retroactive disability payments on recipients' compliance with substance abuse treatment. *Psychiatric Services* 48:796–799, 1997
5. Grossman LS, Willer JK, Miller NS, et al: Temporal patterns of veterans psychiatric service utilization, disability payments, and cocaine use. *Journal of Psychoactive Drugs* 29:285–290, 1997
6. Phillips DP, Christenfeld N, Ryan NM: An increase in the number of deaths in the United States in the first week of the month: an association with substance abuse and other causes of death. *New England Journal of Medicine* 341:93–98, 1999
7. O'Brien CP, Childress AR, McLellan T, et al: Integrating systemic cue exposure with standard treatment in recovering drug dependent patients. *Addictive Behaviors* 15:355–365, 1990
8. Burns BJ, Santos AB: Assertive community treatment: an update of randomized trials. *Psychiatric Services* 46:669–675, 1995
9. Ries RK, Dyck DG: Representative payee practices of community mental health centers in Washington State. *Psychiatric Services* 48:811–814, 1997
10. Chan K, Fitzsimmons V, Hardy R, et al: *All My Money: A Financial Management Curriculum for Persons Working With Resource-Limited Audiences*. Chicago, University of Illinois Trustees, 2005
11. Essock SM, Mueser KT, Drake RE, et al: Comparison of ACT and standard case management for delivering integrated treatment for co-occurring disorders. *Psychiatric Services* 57:185–196, 2006
12. Morse GA, Calsyn RJ, Dean Klinkenberg W, et al: Treating homeless clients with severe mental illness and substance use disorders: costs and outcomes. *Community Mental Health Journal* 42:377–404, 2006
13. Tsemberis S, Gulcur L, Nakae M: Housing first, consumer choice, and harm reduction for homeless individuals with a dual diagnosis. *American Journal of Public Health* 94:651–656, 2004
14. Conrad KJ, Lutz G, Matters MD, et al: Randomized trial of psychiatric care with representative payeeship for persons with serious mental illness. *Psychiatric Services* 57:197–204, 2006
15. Rosen MI, McMahon TJ, Rosenheck R: Does assigning a representative payee reduce substance abuse? *Drug and Alcohol Dependence* 86:115–122, 2007
16. Luchins DJ, Hanrahan P, Conrad KJ, et al: An agency-based representative payee program and improved community tenure of persons with mental illness. *Psychiatric Services* 49:1218–1222, 1998
17. Stout RL, Wirtz PW, Carbonari JP, et al: Ensuring balanced distribution of prognostic factors in treatment outcome research. *Journal of Studies on Alcohol Supplement* 12:70–75, 1994
18. Rosen MI, Bailey M, Rosenheck RR: Alcohol and drug abuse: principles of money management as a therapy for addiction. *Psychiatric Services* 54:171–173, 2003
19. Petry NM: A comprehensive guide to the application of contingency management procedures in clinical settings. *Drug and Alcohol Dependence* 58:9–25, 2000
20. Rosen MI, Rosenheck R, Ablondi K: *ATM: A Clinician Manual to Conduct Money Management With Your Clients*. West Haven, Conn, Yale University, Department of Psychiatry. Available at [www.behaviorchange.yale.edu/resources/docs/ATM-ManualFinal.pdf](http://www.behaviorchange.yale.edu/resources/docs/ATM-ManualFinal.pdf)
21. First MB, Spitzer RL, Gibbon M, et al: *Structured Clinical Interview for Axis I DSM-IV Disorders—(SCID)*. Patient ed, version 2.0. New York, New York State Psychiatric Institute, Biometrics Research Department, 1996
22. McLellan AT, Kushner H, Metzger D, et al: The fifth edition of the Addiction Severity Index. *Journal of Substance Abuse Treatment* 9:199–213, 1992
23. Fals-Stewart W, O'Farrell TJ, Freitas TT, et al: The Timeline Followback reports of psychoactive substance use by drug-abusing patients: psychometric properties. *Journal of Consulting and Clinical Psychology* 68:134–144, 2000
24. Spitzer RL, Gibbon M, Endicott J: Global Assessment Scale (GAS), Global Assessment of Functioning (GAF) Scale, Social and Occupational Functioning Assessment Scale (SOFAS), in *Handbook of Psychiatric Measures*. Edited by Rush AJ, Pincus HA, First MB, et al. Washington, DC, American Psychiatric Association, 2000
25. Rosen MI, Bailey M, Dombrowski E, et al: A comparison of satisfaction with clinician, family members/friends and attorneys as payees. *Community Mental Health Journal* 41:291–306, 2005
26. Alterman AI, Brown LS, Zaballero A, et al: Interviewer severity ratings and composite scores of the ASI: a further look. *Drug and Alcohol Dependence* 34:201–209, 1994
27. Dixon L, Turner J, Krauss N, et al: Case managers' and clients' perspectives on a representative payee program. *Psychiatric Services* 50:781–786, 1999
28. Evans JD, Wright DE, Svanum S, et al: Psychiatric disorder and unmet service needs among welfare clients in a representative payee program. *Community Mental Health Journal* 40:539–548, 2004
29. Maslow AH: *Motivation and Personality*. New York, Harper and Row, 1954
30. Petrakis IL, Poling J, Levinson C, et al: Naltrexone and disulfiram in patients with alcohol dependence and comorbid psychiatric disorders. *Biological Psychiatry* 57:1128–1137, 2005
31. Hsieh FY: Sample size formulae for intervention studies with the cluster as unit of randomization. *Statistics in Medicine* 7:1195–1201, 1988