Diagnosis Category	ICD-9 Codes			
Alcohol and Drug Bsychosic	291.0-291.9			
Alcohol and Drug Psychosis	292.0-292.9			
Schizophrenic psychoses	295.00-295.95			
	296.00-296.06			
Affective provehouses	296.10-296.16			
Affective psychoses	296.20-296.36			
	296.40-296.99			
Paranoid state psychoses	297.0-297.9			
Other nonorganic psychoses	298.0-298.9			
Neurotic Disorders	300.00-300.9			
Personality Disorders	301.0-301.9			
Alcohol dependence	303.0 -303.93			
Drug dependence	304.00-304.99			
Nondependent abuse of drugs	305.00-305.93			
Special symptoms or syndromes	307.1-307.59			
Adult reaction to stress	308.0-308.9			
Adjustment reaction, including PTSD	309.0-309.9			
Depressive disorder	311			
Montal disorders diagnosed in childhood	312.00-312.9			
Mental disorders diagnosed in childhood	314.00-314.9			
<sup>1</sup> After October 1, 2015, manned ICD 10 codes were used				

## ICD-9 Codes of mental health diagnosis inclusion criteria<sup>1</sup>

<sup>1</sup> After October 1, 2015, mapped ICD-10 codes were used.


## CPT Code inclusion list to define a psychotherapy visit

CPT Code	CPT Description				
90791	PSYCHIATRIC DIAGNOSTIC EVALUATION				
90832	PSYCHOTHERAPY, 30 MINUTES WITH PATIENT				
90834	PSYCHOTHERAPY, 45 MINUTES WITH PATIENT				
90837	PSYCHOTHERAPY, 60 MINUTES WITH PATIENT				
90839	PSYCHOTHERAPY FOR CRISIS; IRST 60 MINUTES				
	FAMILY PSYCHOTHERAPY (CONJOINT PSYCHOTHERAPY) (WITH PATIENT PRESENT), 50				
90847	MINUTES				
90849	MULTIPLE-FAMILY GROUP PSYCHOTHERAPY				
90853	GROUP PSYCHOTHERAPY (OTHER THAN OF A MULTIPLE-FAMILY GROUP)				
	PHYSICIAN OR OTHER QUALIFIED HEALTH CARE PROFESSIONAL QUALIFIED BY EDUCATION,				
	TRAINING, LICENSURE/REGULATION (WHEN APPLICABLE) EDUCATIONAL SERVICES				
	RENDERED TO PATIENTS IN A GROUP SETTING (EG, PRENATAL, OBESITY, OR DIABETIC				
99078	INSTRUCTIONS) OBESITY, OR DIABETIC INSTRUCTIONS)				
99509	HOME VISIT FOR ASSISTANCE WITH ACTIVITIES OF DAILY LIVING AND PERSONAL CARE				
99510	HOME VISIT FOR INDIVIDUAL, FAMILY, OR MARRIAGE COUNSELING				
H0004	BEHAVIORAL HEALTH COUNSELING AND THERAPY, PER 15 MINUTES				
H0005	ALCOHOL AND/OR DRUG SERVICES; GROUP COUNSELING BY A CLINICIAN				
	ALCOHOL AND/OR DRUG SERVICES; INTENSIVE OUTPATIENT (TREATMENT PROGRAM THAT				
	OPERATES AT LEAST 3 HOURS/DAY AND AT LEAST 3 DAYS/WEEK AND IS BASED ON AN				
	INDIVIDUALIZED TREATMENT PLAN), INCLUDING ASSESSMENT, COUNSELING; CRISIS				
H0015	INTERVENTION, AND ACTIVITY THERAPIES OR EDUCATION				
	ALCOHOL AND/OR DRUG SERVICES; INTENSIVE OUTPATIENT (TREATMENT PROGRAM THAT OPERATES AT LEAST 3 HOURS/DAY AND AT LEAST 3 DAYS/WEEK AND IS BASED ON AN				
	INDIVIDUALIZED TREATMENT PLAN), INCLUDING ASSESSMENT, COUNSELING; CRISIS				
H0015	INTERVENTION, AND ACTIVITY THERAPIES OR EDUCATION				
H0031	MENTAL HEALTH ASSESSMENT, BY NON-PHYSICIAN				
H0036	COMMUNITY PSYCHIATRIC SUPPORTIVE TREATMENT, FACE-TO-FACE, PER 15 MINUTES				
H2011	CRISIS INTERVENTION SERVICE, PER 15 MINUTES				
H2011	PSYCHOSOCIAL REHABILITATION SERVICES, PER 15 MINUTES				
H2027	PSYCHOEDUCATIONAL SERVICE, PER 15 MINUTES				
H2U27	PATIENT EDUCATION, NOT OTHERWISE CLASSIFIED, NON-PHYSICIAN PROVIDER,				
S9445	INDIVIDUAL, PER SESSION				
00110	PATIENT EDUCATION, NOT OTHERWISE CLASSIFIED, NON-PHYSICIAN PROVIDER, GROUP,				
S9446	PATIENT EDUCATION, NOT OTHERWISE CLASSIFIED, NON-PHYSICIAN PROVIDER, GROUP, PER SESSION				
33440					

## Definition of medication management encounter

Veterans with MH or SUD diagnoses who had at least one E& M-coded visit (CPT codes: 90792, 90863, 99202, 99203, 99204, 99205, 99213, 99214, 99215, 99342, 99343, 99344, 99345, 99348, 99349, 99350, 99442, 99443, 99605, 99606, 99607) with a psychiatrist or at least one E& M-coded visit with another qualifying prescriber in a MH clinic. E& M visits include office visits, home visits and telephone visits. Other prescribers include personclass codes corresponding with non-psychiatrist MDs (Other MH MD), Clinical Pharmacists (Pharmacist), Physician Assistants (MH PA), Clinical Nurse Specialists (MH CNS) or Nurse Practitioners (MH NP) on the Person Code Report. E& M visits with non-psychiatrist prescribers count only if coupled with the following clinic stops: 156, 157, 292, 500 to 503, 505, 506, 509, 510, 512 to 521, 523 to 527, 531 to 542, 545 to 548, 550 to 568, 571 to 584, 586 to 589, 593 to 599, 713.

## Sample Sizes flow diagram for all analyses



Descriptive overview of tablet users vs. non-users (weighted)

	Tablet used Tablet not used		
	Mean/%	Mean/%	р
Age (M±SD)	58.34 ± 14.17	61.07 ± 14.78	<.001
Sex (male=1) (%)	95.27%	93.28%	<.001
Race (%)			
Hispanic	5.30%	5.30% 5.04%	
Non-Hispanic White	79.73%	76.47%	0.002
Non-Hispanic Black	11.55%	15.97%	<.001
Other	3.41%	2.52%	.030
LTE coverage (%)	98.85%	99.26%	<.001
Rurality (%)			
Urban	57.95%	63.87%	.020
Rural	39.96%	34.45%	.065
Highly rural	2.08%	1.68%	.002
Drive distance to primary care (>40 miles)			
(%)	5.11%	6.72%	.269
Drive distance to primary care (M ± SD)	16.39 ± 14.50	15.07 ± 14.08	.138
Chronic conditions (%)†			
Zero	3.79%	3.36%	.479
1 to 2	18.94%	19.33%	.348
3 or more	77.27%	77.31%	.294
No. of baseline psychotherapy visits (M ±	2.44 + 0.00		010
SD) No. 3 month baseline psychotherapy visits	$2.11 \pm 8.00$	1.95 ± 6.75	.010
$(M \pm SD)$	.38 ± .89	.42 ± 1.19	.632
Baseline MH diagnoses (%)	.50 ± .05	.42 ± 1.15	.052
PTSD	9.09%	12.61%	<.001
SUD	46.40%	31.93%	.095
Depression	10.61%	13.45%	.158
SMI	54.73%	52.10%	<.001
New mental health diagnosis (%)	14.20%	11.76%	.864
2 or more mental health diagnoses (%)	35.61%	23.53%	<.001
No. Baseline MH encounters ( $M \pm SD$ )	8.91 ± 14.45	8.02 ±14.35	<.001
SuicideRiskFlag (%)	1.89%	1.68%	<.001
n	583	145	
	505	143	

**†** From list of 34 chronic conditions

LTE = Long Term Evolution, MH = Mental Health, SMI = Serious Mental Illness, PTSD = Post Traumatic Stress Disorder, SUD = Substance Use Disorder, Suicide risk flag = VHA-based algorithm identifying patients at high-risk for suicide

Baseline and follow-up outcomes for tablet users, non-users, and control cohort (weighted)

	Baseline			Follow up		
	Tablet used	Tablet not used	Control	Tablet used	Tablet not used	Control
No. Psychotherapy visits (M ± SD)	0.71 ± 1.68	1.14 ± 5.31	.62 ± 3.14	2.82 ± 6.99	0.79 ± 2.03	.30 ± 1.27
No. Medication management visits (M ± SD)	1.72 ± 2.19	1.12 ± 1.36	1.21 ± 2.15	1.93 ± 2.52	1.24 ± 3.43	.36 ± 1.36
SAIL Measure met (Yes=1) (%)	3.43%	2.07%	3.64%	24.19%	6.21%	2.55%
Missed opportunity rate (M ± SD %)	29.00% ±		30.13% ±	27.30% ±		50.01% ±
	30.10	25.85% ± 30.44	33.35	28.04	30.7% ± 34.10	45.34
Any ED visit (Yes=1) (%)	12.01%	10.34%	8.09%	16.98%	17.93%	11.51%
No. of ED visits (M ± SD)	.31 ± 1.30	.22 ± .93	.21 ± 1.18	.41 ± 1.37	.42 ± 1.18	.27 ± 1.36
n	583	145	1,020	583	145	1,020