

Supplement Table. Frequency of Mental Distress, Comparison between Psychiatric Groups and a Non-psychiatric Group¹

	Log Rank Test Score	Odds Ratios and 95% Confidence Intervals		
		Schizophrenia (N=50)	Bipolar Disorder (N=51)	Major Depressive Disorder (N=29)
1-Nervous	31.31***	2.54 (1.20, 5.35)*	2.75 (1.36, 5.53)**	5.02 (2.20, 11.45)**
2-Depressed	31.11***	3.40 (1.46, 7.61)**	4.30 (1.99, 9.27)***	4.82 (2.09, 11.14)***
3-Hyperarousal	26.03***	6.46 (1.81, 23.06)**	12.16 (3.61, 41.00)***	7.72 (2.05, 29.04)**
4-Lonely	24.98***	4.02 (1.86, 8.69)***	2.79 (1.38, 5.64)**	4.74 (2.03, 11.06)***
5-Hopeful	19.94**	0.45 (0.21, 0.97)*	0.41 (0.21, 0.82)**	0.37 (0.16, 0.85)*
6-Trouble sleeping	19.27**	0.38 (0.17, 0.83)*	1.27 (0.64, 2.52)	2.64 (1.18, 5.89)*

¹ The numbers of the items correspond to the numbers of the items in Figure 1. Items inquired, “In the past week, how often have you ...” Item responses were rated on a 4 point scale from “Rarely” to “Most of the time.” Items were 1-Felt nervous, anxious, or on edge; 2- Felt depressed; 3 -Had physical reactions when thinking about your experience with the COVID-19 epidemic; 4- Felt lonely; 5 -Felt hopeful about the future (direction of responses reversed); 6 -Had trouble sleeping. Odds Ratios and Log Rank Test scores from ordered logistic regression analyses with the non-psychiatric individuals (n=65) as the reference group. All analyses adjusted for age, race, and gender. $p > .05$ except where indicated; * $p < .05$, ** $p < .0125$, *** $p < .001$.

Supplement Table. COVID-related Concerns, Comparison between Psychiatric Groups and a Non-psychiatric Group¹

	Log Rank Test Score	Odds Ratios and 95% Confidence Intervals		
		Schizophrenia (N=50)	Bipolar Disorder (N=51)	Major Depressive Disorder (N=29)
1-Overwhelmed	24.21***	4.36 (2.06, 9.20)***	1.35 (0.68, 2.67)	1.52 (0.69, 3.37)
2-Medical bills	19.69**	3.96 (1.86, 8.43)***	1.95 (0.98, 3.90)	1.12 (0.50, 2.50)
3-Food	22.36***	3.88 (1.77, 8.47)***	2.98 (1.46, 6.09)**	1.39 (0.61, 3.16)
4-Concentration	14.59*	3.41 (1.63, 7.14)***	1.28 (0.63, 2.62)	1.71 (0.77, 3.80)
5-Anxious	12.77*	2.90 (1.40, 6.04)**	1.49 (0.76, 2.94)	1.87 (0.83, 4.25)
6-Money	15.88*	2.36 (1.11, 4.99)*	2.53 (1.28, 4.98)**	1.21 (0.55, 2.69)
7-Infection-Self	16.22*	2.22 (1.06, 4.64)*	0.51 (0.26, 1.01)	0.73 (0.34, 1.59)
8-Income	9.50	1.71(0.82, 3.55)	1.33 (0.68, 2.61)	0.65 (0.29, 1.47)
9-Spending	7.01	1.57 (0.76, 3.26)	1.29 (0.65, 2.57)	1.18 (0.54, 2.60)
10-Sleep	6.56	1.49 (0.71, 3.11)	1.51 (0.74, 3.10)	2.15 (0.95, 4.87)
11-Infection-Others	2.67	1.35 (0.64, 2.82)	1.06 (0.54, 2.11)	0.99 (0.44, 2.24)
12-Transmission	11.81	0.70 (0.34, 1.35)	0.67 (0.34, 1.29)	0.76 (0.35, 1.69)

¹ The numbers of items correspond to the numbers of items in Figure 2. Items were rated on 5 point scale from “Strongly Disagree” to “Strongly Agree.” Items were: 1-I am feeling overwhelmed by COVID-19; 2- I am worried about medical bills if I get sick from COVID-19; 3- I am worried about having enough food because of COVID-19; 4- I have difficulties concentrating because of COVID-19; 5- Thinking about COVID-19 makes me very anxious; 6- I am worried about money because of COVID 19; 7- I am very worried about getting COVID-19;8- I am worried about loss of income if I get sick from COVID-19; 9- I am spending more money because of COVID-19; 10- I have a hard time sleeping because of COVID-19; 11- I am very worried about my family/friends getting COVID-19; 12 -I am very worried about giving someone else COVID-19. Odds Ratios and Log Rank Test scores from ordered logistic regression analyses with the non-psychiatric individuals (n=65) as the reference group. All analyses adjusted for age, race, and gender. p>.05 except where indicated; * p<.05, **p<.0125, ***p<.001.