

## Demographic characteristics and prescribed index antipsychotic medications

Characteristic	N	%
<b>Race</b>		
Black	169	68.9
White	68	27.8
Other	8	3.3
<b>Age</b>		
23-33 years	81	33
34-44 years	83	34
45-65 years	81	33
<b>Gang affiliation</b>		
Yes	32	13.1
No	213	86.9
<b>Index Prescriptions</b>		
Olanzapine	84	34.3
Risperidone	68	27.8
Haloperidol	33	13.5
Chlorpromazine	14	5.7
Haloperidol decanoate	9	3.7
Fluphenazine	7	2.9
>1 medication	7	2.9
refused medication	6	2.4
Fluphenazine decanoate	5	2.0
Perphenazine	4	1.6
Ziprasidone	3	1.2
Aripiprazole	2	0.8
Thiothixene	2	0.8
Trifluoperazine	1	0.4

Rates and relative risk (RR) of adverse events during and after stopping antipsychotic medication

	Rate*		RR	(95% CIs)	p-value
	During	After			
Psychiatric hospitalizations	0.4	0.7	0.63	(0.30, 1.30)	0.208
Emergent suicide assessments	0.9	0.6	1.50	(0.65, 3.42)	0.340
Days on suicide watch	2.8	1.7	1.61	(0.32, 8.21)	0.567
Number of infractions	3.1	5.3	0.58	(0.36, 0.94)	0.024**
Days assigned to DSEG	67.7	96.6	0.68	(0.43, 1.08)	0.102

\*per 1000 person-days  
 Disciplinary segregation  
 (DSEG)  
 Confidence intervals (CIs)  
 \*\* p < 0.05