

**ONLINE ONLY APPENDIX TABLE 2. OFFICE VISITS FOR BIPOLAR DISORDER IN WHICH AN ATYPICAL ANTIPSYCHOTIC WAS USED, 1998-2009.**

Atypical Antipsychotics ...	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
...Used Alone, %	7	11	10	13	10	17	20	20	21	25	23	27
...Used with Mood Stabilizer, %	77	72	71	63	63	61	51	55	55	52	51	47
...Used with Antidepressant, %	38	38	40	40	47	43	43	38	37	35	37	38
...Used with Anxiolytic, %	17	19	15	12	15	14	17	16	17	17	16	17
...Used with Other Therapy, %	4	3	5	5	3	4	3	3	4	3	4	5
<b>Total Visits where atypical used (millions)</b>	<b>0.7</b>	<b>0.7</b>	<b>0.9</b>	<b>1.2</b>	<b>2.0</b>	<b>2.2</b>	<b>2.4</b>	<b>3.3</b>	<b>4.0</b>	<b>4.1</b>	<b>3.9</b>	<b>3.8</b>

Column percents exceed 100% because rows are not mutually exclusive (e.g., atypical prescribed with both antidepressant and anxiolytic would contribute to two rows); "Other" includes stimulants and typical antipsychotics;  
Source: IMS Health National Disease and Therapeutic Index™, 1998-2009

