

Continuity of Care and Discharge Planning for Hospital Psychiatric Admissions

Online Supplement

Unadjusted and adjusted associations between demographic, clinical characteristics and attending an outpatient appointment within 7 days of discharge for the total sample of 189 psychiatric inpatients.							
		Attended outpatient appointment within 7 days following discharge (n=99, 52%)		Unadjusted regressions		Adjusted regression	
		n	% (Row)	OR	95% CI	OR	95% CI
Communication with outpatient provider	No	28	39	ref		ref	
	Yes	71	60	2.32	1.27-4.24	1.65	0.82-3.32
Attended outpatient mental health appointment in the 30 days prior to admission	No	46	44	ref		ref	
	Yes	53	62	2.09	1.16-3.75	1.75	0.88-3.49
Age	Youth (20 and under)	19	61	ref		ref	
	Adult (21+)	80	51	0.65	0.29-1.42	0.71	0.27-1.86
Gender	Female	52	56	ref		ref	
	Male	47	49	0.76	0.43-1.34	0.84	0.44-1.61
Race/ethnicity	Black non-Hispanic	36	43	ref		ref	
	Hispanic/Puerto Rican	12	57	1.74	0.66-4.58	1.38	0.48-3.93
	White non-Hispanic	41	62	2.14	1.11-4.14	1.95	0.95-4.00
	Other/unknown	10	53	1.45	0.53-3.94	1.19	0.40-3.53
Inpatient length of stay	1-6 days	19	45	ref		ref	
	7-13 days	42	55	1.50	0.70-3.19	1.57	0.69-3.56
	14 days or longer	38	54	1.39	0.65-3.00	1.63	0.67-3.95
Primary discharge diagnosis	Psychotic disorder	40	49	ref		ref	
	Mood disorder	49	54	1.25	0.69-2.29	1.19	0.59-2.39
	Other	10	59	1.50	0.52-4.32	1.74	0.49-6.14
Co-occurring substance use disorder	No	57	56	ref		ref	
	Yes	42	48	0.70	0.40-1.25	0.92	0.49-1.74