

Online supplement for DOI 10.1176/appi.ps.202100478

Table 1. Associations between patient population COVID-19 prevalence and the percentage of eligible patients who had depression care follow-up plan documented: 2020.

	Model 1		Model 2	
	Coefficient	SE	Coefficient	SE
Patients diagnosed with COVID-19, %	-0.47*	0.238		
Patients tested for COVID-19, %			-0.10*	0.053
Patient population non-Hispanic black, %	0.15*	0.065	0.16*	0.064
Patient population non-Hispanic white, %	0.21**	0.062	0.22**	0.062
Patient population Hispanic, %	0.21**	0.069	0.21**	0.07
Patient population uninsured, %	0.05	0.052	0.08	0.053
Patient population living below poverty, %	0.06+	0.035	0.07+	0.035
Patient population diagnosed with obesity, %	0.30**	0.037	0.31**	0.038
Total supplemental COVID-19 funding; \$10,000s	-0.002	0.011	-0.002	0.011
Total patients; 1,000s	0.02	0.037	0.02	0.037
State-level COVID-19 cases per capita (cumulative by December 2020)	124.00**	44.954	107.50*	43.33
Constant	30.36**	5.992	29.62**	6.065
Observations	1,113		1,113	
R-squared	0.11		0.12	

Notes: Estimates using data from the UDS and the New York Times. There were $N = 1,113$ observations. SE indicates Huber-White standard errors. + $P < 0.10$, * $P < 0.05$, ** $P < 0.01$.