

Supplement Table 1. Marginal effects of psychotherapy disruption from multilevel models with cross-level interactions for each mental health diagnosis and time period

Mental Health Diagnoses	Marginal Effect with time (During COVID-19 vs. Before COVID-19)		
	aOR	(95% CI)	p-value
ADHD			
Yes	0.53	(0.50, 0.55)	<0.001
No	0.45	(0.44, 0.46)	<0.001
Anxiety Disorder			
Yes	0.47	(0.45, 0.48)	<0.001
No	0.42	(0.41, 0.44)	<0.001
Autism Spectrum Disorder			
Yes	0.57	(0.49, 0.65)	<0.001
No	0.45	(0.44, 0.46)	<0.001
Bipolar Spectrum Disorder			
Yes	0.58	(0.55, 0.62)	<0.001
No	0.44	(0.43, 0.45)	<0.001
Conduct/Disruptive Disorder			
Yes	0.47	(0.39, 0.57)	<0.001
No	0.45	(0.45, 0.46)	<0.001
Dementia			
Yes	0.50	(0.47, 0.54)	<0.001
No	0.45	(0.44, 0.46)	<0.001
Depressive Disorder			
Yes	0.48	(0.47, 0.49)	<0.001
No	0.39	(0.38, 0.40)	<0.001
Personality Disorder			
Yes	0.61	(0.57, 0.66)	<0.001
No	0.45	(0.44, 0.46)	<0.001
Other Psychosis			
Yes	0.54	(0.48, 0.6)	<0.001
No	0.45	(0.44, 0.46)	<0.001
Schizophrenia Spectrum Disorder			
Yes	0.57	(0.51, 0.64)	<0.001
No	0.45	(0.44, 0.46)	<0.001
Substance Abuse Disorder			
Yes	0.49	(0.47, 0.51)	<0.001
No	0.45	(0.44, 0.46)	<0.001

The estimates here represent the effect of "During COVID-19" compared to "Before COVID-19" with adjustment for number of previous psychotherapy visit and sociodemographic characteristics